Preprint typeset using LaTeX style emulateapj v. 03/07/07

arXiv:0709.4069v1 [astro-ph] 26 Sep 2007

LARGE AMOUNTS OF OPTICALLY-OBSCURED STAR FORMATION IN THE HOST GALAXIES OF SOME TYPE-2 QUASARS.

M. Lacy[1] A. Sajina[1], A.O. Petric[1,2], N. Seymour[1], G. Canalizo[3], S.E. Ridgway[4] L. Armus[1], L.J. Storrie-Lombardi [1]

ABSTRACT

We present *Hubble Space Telescope* images, and spectral energy distributions from optical to infrared wavelengths for a sample of six $0.3 < z < 0.8$ type-2 quasars selected in the mid-infrared using data from the *Spitzer Space Telescope*. All the host galaxies show some signs of disturbance. Most seem to possess dusty, star-forming disks. The disk inclination, estimated from the axial ratio of the hosts, correlates with the depth of the silicate feature in the mid-infrared spectra, implying that at least some of the reddening towards the AGN arises in the host galaxy. The star formation rates in these objects, as inferred from the strengths of the PAH features and far-infrared continuum, range from $3-90 M_{\odot}\mathrm{yr}^{-1}$, but are mostly much larger than those inferred from the [OII]3727 emission line luminosity, due to obscuration. Taken together with studies of type-2 quasar hosts from samples selected in the optical and X-ray, this is consistent with previous suggestions that two types of extinction processes operate within the type-2 quasar population, namely a component due to the dusty torus in the immediate environment of the AGN, and a more extended component due to a dusty, star forming disk.

Subject headings: quasars:general – galaxies:Seyfert – infrared:galaxies – galaxies:starburst

1. INTRODUCTION

Co-evolution of black holes and their host galaxies is a prediction of most models for the observed close correlation between black hole mass and bulge luminosity/velocity dispersion in local galaxies (e.g. Kauffmann & Haehnelt 2000; di Matteo, Springel & Herquist 2005). Studies of local AGN and type-1 (unobscured) quasars show a correlation between AGN luminosity in the mid-infrared and starburst luminosity, estimated from the far-infrared emission and/or PAH features (Schweitzer et al. 2006; Shi et al. 2007; Maiolino et al. 2007), and the host galaxies of quasars with strong far-infrared excesses do seem to be forming stars (Canalizo & Stockton 2001). However, in general, host galaxy imaging studies in the optical and near-IR do not suggest that strong star formation occurs simultaneously with quasar activity in the majority of quasars. Imaging of type-1 quasar host galaxies show that the hosts of luminous quasars tend to be elliptical galaxies with little obvious signs of disturbance in *Hubble Space Telescope* (HST) images (e.g. Floyd et al. 2004), and only about 30% of quasars show signs of mergers or interactions in HST or adaptive optics images (Marble et al. 2003; Guyon et al. 2007), though deeper (several orbit) *HST*/ACS imaging often reveals faint tidal features in quasar hosts (Canalizo et al. 2007; Bennert et al., in prepraration). Studies of type-2 (obscured) quasars have so far been consistent with this picture, and with the idea that the obscuration takes place in a nuclear torus. Sturm et al. (2006), in their *Spitzer* study of X-ray selected type-2 quasars, make the claim that "type-2 quasars are not ULIRGs" based on featureless mid-infrared spectra. *HST* imaging studies of the host galaxies of type-2 quasars selected from the Sloan Digital Sky Survey (SDSS) show that they are consistent with being hosted by relatively undisturbed elliptical galaxies, once the likely effects of scattered light are accounted for (Zakamska et al. 2006).

In this Letter, we present results of *Spitzer* spectroscopy and photometry, and *Hubble Space Telescope* (*HST*) imaging of a sample of type-2 quasars, and discuss the implications for the co-evolution of galaxies and black holes. Our sample is selected in the mid-infrared, which is less sensitive to reddening in the host galaxy than X-ray or optical selection, and their type-2 nature allows us to image the central regions of the host with high fidelity. The combination of infrared SEDs and optical imaging of this sample thus allows us for the first time to present a coherent picture of the nature of the star formation in the host galaxies of type-2 quasars.

2. OBSERVATIONS

We selected all six candidate type-2 quasars with $0.3 < z < 0.8$ from the 1mJy 8μm flux density limited sample of Lacy et al. (2004). The redshift range ensures that these objects are luminous enough to be on or above the quasar/Seyfert divide in the mid-infrared (5μm luminosity, $L_{5\mu\mathrm{m}} \sim 10^{23.6}\mathrm{WHz}^{-1}$, Lacy et al. 2005a), but low enough for rest-frame B-band to be within the F814W filter of the Advanced Camera for Surveys (ACS) aboard *HST*. The objects were observed with the Multiband Imaging Photometer (MIPS) and Infrared Spectrograph (IRS) as part of *Spitzer* program 20083, and with HST as program GO-10848 (Table 1). Redshifts were obtained, and optical spectral classifications attempted (Lacy et al. 2007). Low dispersion spectra were also obtained with the Spex instrument (Rayner et al. 2003) in prism mode at the Infrared Telescope Facility (IRTF).

[1] Spitzer Science Center, Caltech, Mail Code 220-6, Pasadena, CA 91125; mlacy@ipac.caltech.edu,sajina@ipac.caltech.edu, lisa@ipac.caltech.edu
[2] Columbia University of New York, NY, USA; andreea@ipac.caltech.edu
[3] University of California, Riverside, CA, USA; gabriela.canalizo@ucr.edu
[4] NOAO, Colina El Pino s/n, Casialla 603, La Serena, Chile; seridgway@ctio.noao.edu

TABLE 1
OBSERVATIONS

Name	IRS reqkey	MIPS reqkey	ACS filters
SSTXFLS J172123.1+601214	14016768	14018304	F625W,F435W
SSTXFLS J171147.4+585839	14016000	14017536	F814W,F555W
SSTXFLS J171324.2+585549	14016256	14017792	F775W,F475W
SSTXFLS J171831.5+595317	14016512	14018048	F775W,F475W
SSTXFLS J171106.8+590436	14015744	14017280	F775W,F475W
SSTXFLS J172458.3+591545	14017024	14018560	F775W,F475W

The IRS data were obtained in stare mode, using the short-low and long-low modules. The IRS short-low data had a background image subtracted, made by taking a median of all the short-low observations in the Astronomical Observation Request. The long-low data were combined by subtracting nodded pairs, then combining the pairs with minmax rejection. Extraction was performed using the SPICE tool[5], with optimal extraction. Aperture photometry was performed on the pipeline (post-BCD) MIPS mosaics using standard aperture corrections[6].

The ACS data were taken in two filters, one longward and one shortward of the 4000Å break in the rest frame of the galaxy. Observations consisted of four exposures in each filter in a standard dither pattern, for one orbit per object per filter. The ACS pipeline mosaics were also adequate for the analysis performed here. A full analysis of the data will be presented in Seymour et al. (2007, in preparation).

Four of our six objects were easily classified as type-2 quasars in Lacy et al. (2007) based on well established optical emission line diagnostics. However, SSTXLFS J171106.8+590436 and SSTXFLS J172458.3+591525 had optical emission lines properties which were ambiguous. For SSTXFLS172458.3+591525, we are able to estimate a ([SII]6719+6732)/(Hα+[NII]) ratio of 0.4 from the IRTF/Spex spectrum, which, together with an [OIII]/Hβ ratio of 10 and assuming typical [NII]6584/Hα ratios of 0.1-1.6, places this object firmly amongst the AGN in the diagnostic plot of Kewley et al. (2006), with a range in $\log([\mathrm{SII}]/\mathrm{H}\alpha) \approx -0.4 - 0.1$. In the case of SSTXFLS 171106.8+590436, the Hα and [SII] lines are too weak to do this. However, the very high [OIII]/Hβ ratio of $> 10:1$, detection of HeII4686, and a marginal detection of [NeV]14.3μm in the IRS spectrum make it very likely that an AGN classification for the optical spectrum is correct. We thus assume for the rest of this paper that all six objects contain powerful AGN.

3. ANALYSIS

We fit our SEDs and spectra using a model based on those of Sajina et al. (2005, 2006). The IRS spectra were fit directly, along with photometric points in r, i and z bands from the SDSS, IRAC 3.6 and 4.5μm data (Lacy et al. 2005b), MIPS 70 and 160μm data (Frayer et al. 2006 and this paper) and near-infrared photometry from IRTF. We also checked the IRS spectrophotometry against the IRAC 5.8 and 8.0μm and MIPS 24μm (Fadda et al. 2006) broad-band photometry, and found them to agree within $\approx 10\%$.

[5] http://ssc.spitzer.caltech.edu/postbcd/spice.html
[6] http://ssc.spitzer.caltech.edu/mips/apercorr/

The far-infrared (FIR) emission was fit as a modified black body with temperature $\approx$ 45K. A warm (small grain) component was included with a power-law index of ≈ 2, and cutoffs at high and low frequency. The FIR luminosity, L_{FIR}, was estimated by summing the warm and cold components. The hot dust component was based on a power-law with variable spectral index. The long wavelength cutoff was determined by a fit of a Fermi function to the least starforming AGN in our sample, SSTXLFS J172123.1+601214, which shows a turndown in the hot dust emission beginning at a rest-frame wavelength $\approx 20\mu$m. Netzer et al. (2007) also suggest that the AGN-related hot dust emission starts to decrease beyond $\approx 20\mu$m in type-1 quasars. The hot dust emission was exponentially cutoff at high frequencies with a variable cutoff frequency. The unreddened luminosity of the AGN, L_{AGN}, was estimated by integrating this component. The hot dust component is then reddened by the Galactic Center extinction law of Chiar & Tielens (2006), likely to be a good approximation to the extreme density environments of AGN (Sajina et al. 2007).

A modified version of the starburst Polycyclic Aromatic Hydrocarbon (PAH) emission template of Sajina et al. (2007) was used to fit the PAH features. AGN are known to modify the PAH spectra (Smith et al. 2007). In particular, for our objects, we found a better match by boosting the 7.7μm feature in the template by hand. This PAH template was fitted to the SEDs of all galaxies except SSTXFLS J172123.1+601214, which had a small amount of silicate emission added instead. Total PAH luminosities, L_{PAH}, were determined by integrating over the fitted template. Finally, a 5Gyr old single stellar population from Bruzual & Charlot (2003) was fit to the optical points. The basic properties of the fits are listed in Table 2. Parameters which are not well-constrained by the data (for example the warm dust continuum strength in objects not detected at 160μm) are held fixed. Full details of the fitting process will be given in a future paper (Ridgway et al. 2007, in preparation).

4. DISCUSSION

We first address the amount of star formation in the host galaxies. Star formation rates (SFRs) derived from FIR emission alone are suspect for quasars, as reprocessed quasar emission can, in principle, contribute to the FIR flux. In the 5/6 cases where PAHs are detected, however, they show typical ratios of PAH/FIR luminosities over a wide range of L_{FIR} (0.02-0.06 in total PAH integrated over our template, corresponding to 0.0014-0.0042 in the 6.2μm feature, within the range for star-forming galaxies and ULIRGs seen by Peeters, Spoon & Tielens [2004] of $\sim 0.005 - 0.1$). This indicates that the bulk of the FIR emission is indeed likely to be from star formation (see also Sajina et al., in preparation). A similar result has been found for type-1 quasars by Schweitzer et al. (2006). As Figure 2 shows, however, the [OII]3727 emission (from the spectra of Lacy et al. [2007], aperture corrected using the SDSS photometry) is significantly below that expected in an unreddened starburst in most cases, particularly for the objects with high FIR luminosities. This is consistent with the result of Ho (2005), who found low [OII]3727 emission from type-1 quasar hosts compared to their expected star formation rates based on measurements of molecular gas. How-



FIG. 1.— SEDs and HST images of the six type-1 quasars, in order of increasing silicate absorption. The solid black lines are the IRTF/Spex and *Spitzer*/IRS spectra (both smoothed to reduce noise) and the black open circles photometry from SDSS, IRTF and *Spitzer*. The model fit is shown as the red dashed line, with, from left to right, the stellar component, hot dust component (including absorption, but excluding PAH emission) and sum of warm and cold dust components shown as blue chain-dotted lines. To the right of the SEDs we show expanded plots of the IRS spectra around the PAH wavelengths. The ACS images have the long wavelength image colored red and the short wavelength image in green.



FIG. 2.— The ratio of narrow line luminosity in [OII]3727 to FIR luminosity (circles) and in [NeII]12.8μm to FIR luminosity (triangles). Host galaxies dominated by a disk component are shown as filled symbols, SSTXFLS J172123.1+601214, which has an elliptical host, is shown with open symbols. The dashed line is the ratio of [OII] to FIR luminosity expected in an unextincted starburst.



FIG. 3.— Absorption depth in the 9.7μm silicate feature plotted against axial ratio. Symbols as for Figure 2. The dashed lines represent the predictions of simple uniform disk models of differing opacity measured perpendicular to the disk.

ever, the ratio of the [NeII]12.8μm to the FIR luminosity is similar for all our objects. [NeII] is of similar ionization to [OII] and is also therefore dominated by emission from the starburst. This suggests that dust reddening is to blame for the lack of [OII]3727 emission, and is also consistent with the host galaxy colors and morphologies. As can be seen in Figure 1, the host galaxies of the objects with high $L_{\rm FIR}$ are typically disturbed, red objects, likely to be late stage merger remnants. This suggests that star formation in a significant fraction of quasar host galaxies is heavily obscured in the optical.

Obscuration by dusty starforming regions also seems to affect the extinction to the AGN. Figure 3 shows a plot of axial ratio of the host galaxy (measured using the *ellipse* task in IRAF on the longer wavelength image) versus the depth of the 9.7μm silicate feature, $S_{\rm sil}$, as defined by Levenson et al. (2006). Although our sample is small, the five of our six host galaxies with apparent or possible disk-like structures, and 1d-profiles close to exponential disks (excluding morphological disturbances and bulge components), represented by the solid points, seem to show a trend in the sense that the more highly-extincted objects have more edge-on inclinations (lower axial ratios). The one galaxy which does not appear disk-like, SSTXFLS J172123.1+601214 (open symbol), has an underlying $r^{1/4}$ law profile, and is probably an example of an elliptical host merging with a smaller gas-rich galaxy, and lacks the large amount of star formation seen in the mergers of two gas-rich galaxies. In this case, the obscuration is probably almost entirely due to the torus. For the disk-like hosts, a simple uniform dusty disk model appears to work for most, with a spread of about a factor of two in opacity for a line of sight perpendicular to the disk. One object (SSTXLFS J171106.8+590436), however, would have a high opacity in this model even if viewed face-on, corresponding to $S_{\rm sil} \approx -1$. This may be because the extinction is patchy, and a high inclination significantly increases the chance of the line of sight passing through a high opacity region such as a dense molecular cloud. Our two most obscured type-2 quasars (with $S_{\rm sil} = -1.6$ and -2) have silicate absorption as deep or deeper than the most obscured AGN in the sample of Shi et al. (2006). These radio galaxies and Seyfert 2s have $S_{\rm sil} \geq -1.6$, suggesting that extinction can be very extreme in the radio quiet quasar population.

The implication of our result is that many type-2 quasars have at least a component of the extinction towards the nucleus from an extended, star forming disk on scales of kpc, in addition to, or instead of, the traditional obscuring torus, which interferometry suggests is only a few pc in size (Jaffe et al. 2004). Previous suggestions that the host galaxy can contribute significantly to the AGN extinction in type-2 objects have been made by Martinez-Sansigre et al. (2006) and Rigby et al. (2006), based on optical emission line properties. Our results suggest that a significant fraction of the quasar population have their AGN obscured by starforming regions in their host galaxies. Thus studying obscured quasars is essential if we are to understand the origin of the links between the evolution of quasars and their hosts.

We thank the referee for a helpful report. ML and AOP were visiting astronomers at the IRTF, which is operated by the University of Hawaii under Cooperative Agreement no. NCC 5-538 with the National Aeronautics and Space Administration (NASA), Science Mission Directorate, Planetary Astronomy Program. This work is based on observations made with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Support for this work was provided by NASA through an award issued by JPL/Caltech. This work is also based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute (STScI), operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555, and are

TABLE 2
HOST GALAXY PROPERTIES

Object	z	$\log(L_{\rm AGN}$ $[L_\odot])$	$\log(L_{\rm FIR}$ $[L_\odot])$	$\log(L_{\rm PAH}$ $[L_\odot])$	$\log(L_{[\rm OII]}$ $[L_\odot])$	$S_{\rm sil}$	Axial ratio	SFR (FIR) $(M_\odot {\rm yr}^{-1})$	SFR ([OII])† $(M_\odot {\rm yr}^{-1})$
SSTXFLS J172123.1+601214	0.325	11.4	10.6	<9.4	8.17	-	0.63±0.02	3.3	3.9
SSTXLFS J171147.4+585839	0.800	12.1	11.7	10.5	8.44	-0.21	0.96±0.02	56	7.2
SSTXFLS J171324.2+585549	0.609	11.8	11.3	9.8	8.26	-0.34	0.79±0.01	16	4.8
SSTXFLS J171831.5+595317	0.700	11.9	12.0	10.3	8.45	-0.53	0.76±0.05	87	7.4
SSTXFLS J171106.8+590436	0.462	11.8	11.5	10.1	7.49	-1.56	0.67±0.01	33	0.8
SSTXFLS J172458.3+591545	0.494	11.6	11.6	10.4	7.91	-1.97	0.24±0.02	37	2.1

† from Kewley et al. (2004), assuming solar metallicity.

associated with program 10848, partly supported by a grant from STScI.

REFERENCES

Bruzual, G. & Charlot, S. 2003, MNRAS, 344, 1000
Canalizo, G. & Stockton, A.N. 2001, ApJ, 555, 719
Canalizo, G., Bennert, N., Jungwiert, B., Stockton, A., Schweizer, F., Lacy, M. & Peng, C. 2007, ApJ, in press
Chiar, J.E. & Tielens, A.G.G.M. 2006, ApJ, 637, 774
di Matteo, T., Springel, V. & Hernquist, L. 2005, Nat, 433, 604
Fadda, D. et al. 2006, AJ, 131, 2859
Floyd, D., Kukula, M.J., Dunlop, J.S., McLure, R.J., Miller, L., Percival, W.J., Baum, S.A., O'Dea, C.P. 2004, MNRAS, 355, 196
Frayer, D.T. et al. 2006, AJ, 131, 250
Guyon, O., Sanders, D.B. & Stockton, A. 2006, ApJS, 166, 89
Ho, L. ApJ, 2005, 629, 680
Jaffe, W. et al. 2004, Nat, 429, 47
Kauffmann, G. & Haehnelt, M.G. 2000, MNRAS, 311, 576
Kewley, L.J., Geller, M.J. & Jansen, R.A. 2004, AJ, 127, 2002
Kewley, L.J., Groves, B. Kauffmann, G. & Heckman, T. 2006, MNRAS, 372, 961
Lacy, M. et al. 2004, ApJS, 154, 166
Lacy, M. et al. 2005a, MmSAI, 76, 154
Lacy, M. et al. 2005b, ApJS, 161, 41
Lacy, M., Petric, A.O., Sajina, A., Canalizo, G., Storrie-Lombardi, L.J., Armus, L., Fadda, D. & Marleau, F.R. 2007, AJ, 133, 186
Levenson, N.A., Sirocky, M.M., Hao, L., Spoon, H.H.W., Marshall, J.A., Elitzur, M., Houck, J.R. 2007, ApJ, 654, L45
Maiolino, R., Shemmer, O., Imanishi, M., Netzer, H., Oliva, E., Lutz, D. & Sturm, E. 2007, A&A, 468, 979
Marble, A.R., Hines, D.C., Schmidt, G.D., Smith, P.S., Surace, J.A., Armus, L., Cutri, R.M. & Nelson, B.O., 2003, ApJ, 590, 707
Martínez-Sansigre, A. et al. 2006, MNRAS, 370, 1479
Netzer, H. et al. 2007, ApJ, in press, astro-ph/0706.0818
Peeters, E., Spoon, H.W.W. & Tielens, A.G.G.M. 2004, ApJ, 613, 986
Rayner, J.T. et al. 2003, PASP, 115, 362
Rigby, J.R., Rieke, G.H., Donley, J.L, Alonso-Herrero, A., Perez-González, P.G. 2006, ApJ, 645, 115
Sajina, A., Lacy, M. & Scott,D. 2005, ApJ, 621, 256
Sajina, A., Scott, D., Dennefeld, M., Dole, H., Lacy, M. & Lagache, G. 2006, MNRAS, 369, 939
Sajina, A., Yan, L., Armus, L., Choi, P., Fadda, D., Helou, G., Spoon, H. 2007, ApJ, 664, 713
Schweitzer, M. et al. 2006, ApJ, 649, 79
Shi, Y. et al. 2006, ApJ, 653, 127
Smith, J.D.T. et al. 2007, ApJ, 656, 770
Sturm, E., Hasinger, G., Lehmann, I., Mainieri, V., Genzel, R., Lehnert, M.D., Lutz, D. & Tacconi, L.J. 2006, ApJ, 642, 81
Zakamska, N.L. et al. 2006, AJ, 132, 1496